

Chpt 4: Pond Scum to Jet Fuel

this chapter deals with Energy and Cell Metabolism, including the processes of photosynthesis & cellular respiration

we will split this chapter in 2 to cover each of these processes separately

Energy: the capacity to do work

Why do cells need E?

- to make larger more complex molecules

monosacc-----> disacc-----> polysacc

AA + AA+ AA-----> polypeptides

- to break down worn out cell parts

- to move things in & out of cells: active transport

- to move cells: sperm, WBC

- to reproduce, make more cells

Why do organisms need E?

- to grow

by increasing number of cells in body

- to develop

by making more complex structures

cells-----> tissues-----> organs-----> org
sys

- to move in response to environment

- to reproduce

First Law of Thermodynamics or Law of Conservation of E

E of the universe is constant, E cannot be created nor destroyed, instead it can be transferred or transformed

How does this law apply to living organisms?
where do we get our E from?
is E exchange totally efficient?

Second Law of Thermodynamics

as E is transferred, some E is always lost, so
systems tend to change in a way that
increases entropy or disorder
world becomes increasingly disordered as
free E is released

non biological example: my attic

But...life is organized! you started as a single
fertilized egg, you gained cells, tissues,
organs, organ systems
at birth you were much more organized than
that single fertilized egg and that has
continued, you have gotten more complex

how are you defying the 2nd Law?

when do organisms finally obey the 2nd Law?

Cellular Metabolism: the sum total of all the chemical reactions that take place in body
there are 2 kinds of rx: synthesis and decomposition reactions

- synthesis rx (anabolism): make larger molecules

 - what chemical process fits here?

 - in synthesis rx, we put E in and store that E in chemical bonds

 - ex: make carbs fr glucose, prot fr AA

- decomposition rx (catabolism): breaks down molecules

 - what chemical process fits here?

 - in decomposition rx, we release free E

 - ex: break down proteins into AA

what is our food E molecule? glucose

but there is too much E in glucose, so we
convert it to a chemical E molecule

ATP is our chemical E molecule

it is a nucleotide: made of adenine, ribose
& 3 PO₄ groups

ATP: adenosine triphosphate

ADP: adenosine diphosphate

AMP: adenosine monophosphate

A-P~P~P the P to P bonds are high E bonds
when we break these, we release E into the
cell to do work

A-P~P~P----->A-P~P + P + E

ATP----->ADP + P + E releases E

ATP <-----ADP + P + E stores E

In living organisms, E transfers involve Oxidation/ Reduction Reactions

oxidation: the removal of e^- from molecule
those electrons are then given to another

reduction: the addition of e^- to a molecule
these are paired reactions

when you remove e^- from a molecule, it is
lower in E

when you add e^- to molecule, it is higher in E
it is called reduction b/c you are lowering
the charge, thus, making it more negative

OIL RIG: oxidation is loss, reduction is gain
this refers to E

Catalysts: anything that speeds up chemical rx
anything that lowers the activation E
what is activation E?

E required to start a rx
what did you use in chemistry? heat

living things are constantly undergoing
chemical reactions (metabolism)
every rx needs activation E, why can't we
use heat?

Enzymes are biological catalysts
enzymes are proteins
every chem rx in body is catalyzed by a
specific enzyme
names of enzymes end in -ase
ex. lactase

Enzymes are specific

1-each enzyme works on specific chem rx
lactase is enzyme to break down lactose
sucrase breaks down sucrose

2-each enzyme works best in narrow
temperature range
too high temp or too low temp denatures
enzyme
enzyme work best at normal body temp
high fever--enzymes don't work well

3-each enzyme works best in narrow pH range
usually near neutral
high acid denatures enzyme
exception is in the stomach
stomach has pH 2, enzymes here
work best at this pH

denaturing an enzyme: this means the 3D structure of the enzyme is destroyed
remember back to proteins

primary structure: sequence of AA

secondary structure: due to polar R groups, the polypeptide chain begins bending or folding

tertiary structure: due to secondary bending, now other polar parts are attracted and more folding occurs, giving the protein its 3D shape

proteins are not functional until they achieve a 3D shape

too high heat, acid can denature enzymes or destroy the 3D shape, then they no longer function

How do enzymes work?

enzymes have active site: part of enzyme
that matches the starting molecules
starting molecules called substrate

there are 2 models of how enzymes work

1-lock & key

active site of enzyme perfectly fits the
substrates like a key in a lock

2-induced fit

substrates are attracted to active site &
then enzyme changes shape to bring
substrates close together

enzymes do not directly enter rx, they bring
substrates together, then release them,
then enzyme is used again

